

# Emergency Medicine Rapid Reference: Critical Care and Acute Disease Management

MixCache.com

---

## Table of Contents

- **Introduction**
- **Chapter 1** Resuscitation Bay Readiness and the ABCDE Approach
- **Chapter 2** Airway Management: Basic to Difficult Airway Algorithms
- **Chapter 3** Breathing and Mechanical Ventilation: Oxygenation, NIPPV, and Vent Modes
- **Chapter 4** Circulation and Shock: Hemodynamics, Pressors, and Resuscitative Fluids
- **Chapter 5** Sepsis and Septic Shock: Bundles and Time-Zero Care
- **Chapter 6** Cardiac Arrest: High-Performance ACLS and Post-ROSC Care
- **Chapter 7** Dysrhythmias: Recognition and Electrical/Chemical Management
- **Chapter 8** Acute Coronary Syndromes: ED Diagnosis and Reperfusion Pathways
- **Chapter 9** Stroke and Neuroemergencies: FAST Decisions and Thrombolysis/Thrombectomy
- **Chapter 10** Seizures and Status Epilepticus: Rapid Control and Workup
- **Chapter 11** Trauma: Primary and Secondary Survey with Damage Control
- **Chapter 12** Traumatic Brain and Spinal Cord Injury: ICP, MAP, and Neuroprotection
- **Chapter 13** Thoracic Trauma: Air, Blood, and the Failing Chest
- **Chapter 14** Abdominal and Pelvic Trauma: Hemorrhage Control and FAST-Guided Decisions
- **Chapter 15** Orthopedic Emergencies: Reductions, Compartments, and Open Fractures
- **Chapter 16** Burns and Inhalation Injury: Airway, Fluids, and Transfer Criteria
- **Chapter 17** Toxicology I: Approach, Antidotes, and Decontamination
- **Chapter 18** Toxicology II: Common Overdoses and Withdrawal Syndromes
- **Chapter 19** Environmental and Exposure Emergencies: Heat, Cold, Drowning, and Envenomation
- **Chapter 20** Endocrine Crises: DKA, HHS, Adrenal and Thyroid Storm
- **Chapter 21** Renal and Electrolyte Emergencies: Hyperkalemia, Hyponatremia, and AKI
- **Chapter 22** Hematologic and Oncologic Emergencies: DIC, TLS, and Transfusion Reactions
- **Chapter 23** Obstetric and Gynecologic Emergencies: Preeclampsia, Hemorrhage, and Ectopic
- **Chapter 24** Pediatric Resuscitation: PALS, Dosing, and Age-Specific Pitfalls

## Introduction

Emergency care lives in the space between seconds and certainty. This book was created to help clinicians convert that urgency into decisive, lifesaving action. It distills complex critical care and acute disease management into concise, high-yield protocols tailored for the realities of the emergency department—noisy rooms, incomplete histories, shifting diagnoses, and limited time. Every chapter is engineered to move you rapidly from problem recognition to stabilization and definitive next steps.

The intended audience includes emergency physicians, advanced practice providers, residents, medical students, nurses, and prehospital professionals who require quick, actionable guidance at the point of care. Whether you are running a resuscitation bay, staffing a community ED overnight, or initiating prehospital interventions en route, the same principles apply: prioritize threats to life, anticipate clinical turns, and communicate with clarity. Throughout, we emphasize interoperability between prehospital and in-hospital teams so that critical information and momentum are never lost in transition.

Our approach is protocol-driven yet patient-centered. Each topic begins with immediate actions—what to do in the first minutes—followed by focused assessments, key diagnostics, and treatment pathways. Flowcharts and decision aids are designed to clarify rather than constrain, highlighting red flags, branching points, and common pitfalls. Where time matters most—airway crises, shock, sepsis, cardiopulmonary arrest, stroke, trauma, toxic exposures—we foreground the steps that most reliably change outcomes.

Evidence informs our recommendations, but usability governs the format. Protocols reflect current consensus from major guidelines and critical care literature, adapted to the ED context and the realities of variable resources. We point out where equipoise exists, offer pragmatic options when ideal tools are unavailable, and stress frequent reassessment as physiology and information evolve. The goal is not to replace clinical judgment but to scaffold it under pressure.

Human factors are integral to high-quality emergency care. Expert medicine in crisis demands clear role assignment, closed-loop communication, and cognitive offloading. You will find tips on anticipating failure points, leveraging checklists, and using point-of-care diagnostics to shorten time to decision. Pearls and pitfalls embedded in each chapter aim to reduce error, especially at transitions of care and during parallel

tasking.

Finally, this is a rapid reference meant to be carried, marked up, and used. Keep it open beside the monitor, hand it to a teammate to call out steps during a code, and adapt the algorithms to your local pathways. If, in a critical moment, these pages help you recognize a subtle decompensation earlier, deliver a key intervention faster, or coordinate your team more effectively, then the book has done its job—bringing order to urgency and turning seconds into survival.

---

## **Chapter One: Resuscitation Bay Readiness and the ABCDE Approach**

The emergency department is a theater of controlled chaos, a place where minutes often dictate outcomes and every decision carries significant weight. When a critically ill or injured patient rolls through those doors, the initial moments are paramount. This isn't the time for fumbling with equipment or debating who's in charge. It's a time for orchestrated action, built on preparation and a systematic approach. The resuscitation bay, therefore, isn't just a room; it's a meticulously organized cockpit, and your team, a well-drilled flight crew.

Effective emergency care begins long before the ambulance siren wails its arrival. It starts with readiness. Think of it as pre-gaming for the most important competition of your life, every single shift. This proactive stance ensures that when the adrenaline surges, your cognitive load is minimized, allowing you to focus on the patient's immediate needs rather than searching for a laryngoscope blade. It's about building a system that anticipates potential disasters and mitigates them before they even have a chance to unfold.

One of the foundational pillars of this readiness is the "Zero Point Survey." This isn't a complex ritual; it's a simple, yet powerful, mental and physical checklist that takes mere moments but can save precious minutes during a resuscitation. Before the patient even arrives, survey yourself, your team, and your environment. Are you physically and cognitively prepared? Have you addressed your own basic needs like hunger or a full bladder, which, surprisingly, can be significant distractions during high-stakes events?

Next, assess your team. Is there a clearly designated team leader? In the maelstrom of a critical resuscitation, a single, authoritative voice is crucial for preventing confusion and ensuring efficient task allocation. Everyone on the team should know their role, their responsibilities, and to whom they report. Role assignments can be

fluid, but at any given moment, the chain of command should be clear. Briefing the team on the incoming patient, outlining potential life threats, and setting initial priorities also falls under this crucial preparation.

Finally, scrutinize your environment – the resuscitation bay itself. Is all essential equipment present, functional, and easily accessible? This means having advanced airway equipment at the bedside, suction ready, oxygen delivery devices prepped, and intravenous access supplies laid out. Consider setting up for anticipated interventions; for instance, if trauma is expected, having bilateral chest tube kits or a pelvic binder readily available can dramatically reduce time to intervention. The ultrasound machine should be turned on and ready to scan. Medications like tranexamic acid, rapid sequence intubation (RSI) drugs, analgesia, and antiemetics should be immediately at hand. This meticulous preparation of the resuscitation bay ensures that when the patient arrives, the focus shifts entirely to their care, without the added stress of a disorganized environment.

Once the patient arrives and the initial chaos begins to subside, a systematic approach to assessment and treatment becomes your north star: the ABCDE approach. This universally accepted framework, first introduced in parts for cardiopulmonary resuscitation and later expanded for trauma patients, provides a structured method for evaluating and managing critically ill or injured individuals. It's designed to help you quickly identify and address life-threatening conditions in a prioritized order, regardless of the underlying cause.

The ABCDE mnemonic stands for Airway, Breathing, Circulation, Disability, and Exposure. The beauty of this approach lies in its inherent prioritization: you address life-threatening problems in one category before moving on to the next. For example, a compromised airway takes precedence over significant bleeding, because without an open airway, nothing else matters. This systematic process ensures that the most immediate threats to life are identified and managed first, buying precious time for further diagnosis and definitive treatment.

Let's break down each component, understanding that while we discuss them sequentially, many of these assessments and interventions will occur simultaneously in a real-world resuscitation, with different team members tackling different aspects. The overarching goal is to stabilize failing vital functions and transform a complex, overwhelming situation into a series of smaller, manageable problems.

**A: Airway** is always your first priority. Is the patient's airway patent? Can they speak in full sentences? If not, listen for noisy breathing—stridor, gurgling, snoring—all ominous signs of an obstructed or partially obstructed airway. Look for accessory muscle use or paradoxical chest wall movement. A patient who is unconscious or has collapsed should immediately prompt an assessment for airway patency.

If the airway is compromised, immediate intervention is required. This might involve simple maneuvers like a head-tilt/chin-lift or jaw thrust to relieve obstruction caused by the tongue. Suctioning foreign material, blood, or vomit is also critical. If these basic measures aren't sufficient, you'll need to rapidly escalate to advanced airway techniques. This could range from inserting an oropharyngeal or nasopharyngeal airway to, if necessary, definitive airway management like endotracheal intubation. Remember, a secured airway is the foundation upon which all other resuscitative efforts are built.

**B: Breathing** comes next. Once the airway is patent, you need to ensure effective ventilation and oxygenation. Look, listen, and feel. Observe the patient's respiratory rate, depth, and symmetry of chest wall movement. Listen for breath sounds—are they present bilaterally? Are there adventitious sounds like wheezing or crackles? Feel for crepitus or tracheal deviation. Is the patient cyanotic? Are they struggling to breathe?

Immediate interventions for breathing problems often involve providing supplemental oxygen. For patients with respiratory distress, consider non-invasive positive pressure ventilation (NIPPV) like CPAP or BiPAP, if appropriate and tolerated. If breathing is absent or inadequate, bag-valve-mask ventilation is indicated, followed by definitive airway management and mechanical ventilation. Remember, the goal here is not just to move air, but to ensure adequate oxygen delivery to the tissues and carbon dioxide removal.

**C: Circulation** is the third pillar. After ensuring a patent airway and adequate breathing, you need to assess the patient's circulatory status and address any life-threatening issues like hemorrhage or shock. Check for central and peripheral pulses, assessing their presence, rate, quality, regularity, and equality. Weak or thready central pulses, or absent peripheral pulses, are red flags. Measure blood pressure; however, be aware that compensatory mechanisms can initially maintain a normal blood pressure even in shock. A low diastolic pressure can hint at arterial vasodilation, as seen in sepsis or anaphylaxis.

Look for signs of shock: pallor, cool extremities, delayed capillary refill, and altered mental status. Identify and control any obvious external bleeding. Establish intravenous access promptly – ideally two large-bore IVs – and consider initiating fluid resuscitation with crystalloids, or blood products if hemorrhage is suspected. Rapid infusers and warming devices should be employed for large-volume resuscitation. Continuously monitor cardiac rhythm, and if signs of cardiac arrest are present, immediately transition to high-performance cardiopulmonary resuscitation (CPR) protocols.

**D: Disability** refers to a rapid neurological assessment. While a comprehensive

neurological exam takes time, the initial focus is on identifying immediate threats to brain function. The Glasgow Coma Scale (GCS) is a quick and reliable tool to assess a patient's level of consciousness. A rapidly deteriorating GCS demands immediate attention. Assess pupillary size and reactivity to light. Look for gross motor deficits or abnormal posturing.

Hypoxia, hypoglycemia, and hypoperfusion are common reversible causes of altered mental status in the emergency setting. Therefore, ensure adequate oxygenation, check a rapid point-of-care glucose, and address any circulatory compromise. If trauma is involved, assume a cervical spine injury until proven otherwise and maintain spinal precautions.

**E: Exposure** is the final step, but by no means the least important. This involves fully exposing the patient to conduct a thorough head-to-toe examination. This is where you look for injuries or findings that might have been missed in the initial, rapid survey. Carefully inspect the skin for rashes, petechiae, or track marks. Logroll the patient to examine their back for occult trauma or pressure ulcers.

Crucially, after exposing the patient, it's vital to cover them with warm blankets to prevent hypothermia. Critically ill patients are highly susceptible to heat loss, and hypothermia can worsen outcomes, particularly in trauma and sepsis. Maintain patient privacy and dignity throughout this process.

The ABCDE approach is not a one-and-done assessment. It's an iterative process. After each intervention, or at regular intervals, you must reassess the patient, moving through the ABCDEs again to evaluate the effect of your treatments and identify any new or worsening problems. The patient's condition can change rapidly in the ED, and constant vigilance is the hallmark of excellent emergency care.

Beyond the clinical steps, effective communication is paramount. Use clear, concise language. Employ closed-loop communication to confirm that orders are heard, understood, and executed. The SBAR (Situation, Background, Assessment, Recommendation) or RSVP (Reason, Story, Vital signs, Plan) approaches can be incredibly helpful for structured handovers or communicating critical information to consultants. In the heat of the moment, shouting or ambiguous statements only sow confusion. Your team is relying on you for clarity and direction.

Ultimately, the resuscitation bay is where emergency medicine truly shines. It's a crucible where preparedness meets crisis, and a systematic approach transforms potential tragedy into a triumph of timely intervention. By mastering resuscitation bay readiness and flawlessly executing the ABCDE approach, you equip yourself and your team with the tools to confront the most formidable challenges, ensuring that every second counts and every intervention makes a difference.

*This is a sample preview. Purchase the book to read the full content.*

Visit [MixCache.com](http://MixCache.com) to purchase the complete book.